

Integrative Approaches to Gestational Diabetes

Diane Downing
Integrative Family Medicine Fellow
University of Arizona
Program in Integrative Medicine
February 22, 2006



Objectives

- Participants will:
 - understand the usefulness and efficacy of 3 botanicals/supplements used in Diabetes Mellitus and its appropriateness in Gestational Diabetes
 - be familiar with 2 integrative approaches to Gestational Diabetes

Gestational Diabetes

- Defined as a carbohydrate intolerance of variable severity with onset or first recognition during pregnancy



Prevalence-

- Diabetes - most common medical complication of pregnancy
- Incidence of diabetes complicating pregnancy has increased about 40 percent between 1989 and 2002 (Martin and colleagues, 2003).
- In 2002, ~ 131,000 pregnancies complicated by diabetes, representing 3.3 percent of all live births (Martin and colleagues, 2003).
- More than 90 percent were gestational diabetes

Consequences

- More than half of women with gestational diabetes will go on to develop overt diabetes in the ensuing 20 years
- Growing evidence for long range complications in their children that include obesity and diabetes (Feig and Palda, 2002)

Potential Fetal Complications



- Gestational Diabetes-Class A1
 - macrosomia
 - shoulder dystocia
 - brachial plexus injury
 - Neonatal hypoglycemia
 - fetal anomalies are NOT increased (Sheffield and colleagues, 2002)
- Overt Diabetes and Class A2 and above
 - spontaneous abortion
 - preterm delivery
 - stillbirth
 - congenital malformations
 - neonatal morbidity
 - complications as in gestational diabetes

Potential Maternal Complications



■ Gestational

- C-section
- increased risk of CV complications
- elevated lipids
- HTN
- metabolic syndrome

■ Overt

- 10 fold increase risk of maternal death (Cousins, 1987).
- ketoacidosis
- preeclampsia
- retinopathy
- neuropathy
- infections

Goals of therapy

- Fasting plasma glucose < 95 mg/dL
- 2 hour postprandial blood glucose < 120 mg/dL



Mary's Story



Mary

- 26 y/o Hispanic G1P0
- 25 weeks gestation
- 1 hour glucose tolerance test 160 mg/dL
- 3 hour glucose tolerance test also abnormal
- Family Hx – Mom with diabetes mellitus on insulin w/ h/o gestational diabetes when pregnant with Mary
- Eats a diet high in sat fat and carbohydrates

Current Care Strategies

- Diet –
- Exercise –
- Insulin- started when FBG persists >105 mg/dL
- Oral hypoglycemics – not currently recommended by ACOG

Diet



- Nutrition counseling
- 30 kcal/kg/d based on pre-pregnant weight
- if obese – caloric restriction
- Fresh fruits and vegetables
- Low glycemic index carbohydrates
- Whole grains
- High fiber - oats
- Low saturated fat
- Low fat dairy
- Avoid refined sugar

Diet -

■ Oats –

- Pilot crossover study 8 Type II diabetic males
- High fiber oat bran concentrate bread product x 12 weeks (34 gm/d fiber)
- vs. diet with 19 gm/d fiber
- after 24 weeks lower postprandial glucose and insulin levels
- lower total and LDL cholesterol and TGA
- (Pick, et al. 1996)

Diet -

■ Fiber –

- Small randomized crossover study of 13 Type II diabetic patients
- 6 weeks high fiber – 50 gm/d (25 gm soluble + 25 gm insoluble)
- vs. 6 weeks moderate fiber – 24 gm/d (8 gm soluble + 16 gm insoluble)
- Results – decreased preprandial glucose levels by 13 mg/dL, decreased TC, VLDL, TGA (Chandalia, et al. 2000)

Diet –

■ Cinnamon –

- 2003 study – 60 Type II diabetics
- Dose -1,3 or 6 gm/day cassia cinnamon x 40 days
- Results – decreased FBG (18-29%), TGA (23-30%), TC (7-27%) LDL (12-26%) (Khan, et al. 2003)
- 1 tsp. cinnamon = 4.75 grams

Exercise



- Physical activity during pregnancy reduces the risk of gestational diabetes (Dempsey and colleagues 2004)
- Resistance exercise helped avoid insulin therapy in overweight women with gestational diabetes (Brankston and associates 2004)

Potential Herbal Therapy in Diabetes Mellitus

- Gymnema
- Bitter melon
- Prickly Pear
- Fenugreek



Gymnema sylvestre

- Herb used in Ayurveda – a traditional East Indian healing system
- Two non-randomized controlled clinical trials by same research group of Type I and Type II diabetics (Baskaran, et al. 1990. Shanmugasundaram, et al. 1990)
- Showed decreased fasting blood glucose, HgA1C and medication requirements
- Dose – 400 mg GS4 water-soluble leaf extract
- Insufficient data for use in pregnancy



Bitter melon- *Momordica charantia*

- Vegetable eaten regularly in India, Asia, South America and Africa
- Small, preliminary study in Type I and II diabetics using injectable juice extracts showed acute reduction in BG at 30 min, 4 hr and 12 hr. (Baldwa et al. 1977)
- A case series looking at GTT after 50ml bitter melon fruit juice, and then after 8-11 weeks of fried melon bits
- Results - 12% drop in GTT after juice, 6% after melon bits, 8% drop in HgA1c after bits (Leatherdale et al. 1981)



Bitter melon-

- Type I and II diabetics have shown improved glucose tolerance and reduction in mean blood glucose levels (Yeh, et al. 2003)
- Studies of poor quality and small size
- No studies in pregnancy

Bitter Melon - Safety

- Animal study showed a glycoprotein found in seeds to inhibit pregnancy in rats (Chan WY, Tam PP, Yeung HW. 1984.)
- Aqueous *root* extract abortifacient effects in mice (Aguwa CN, Mittal GC. 1983)
- Likely safe when eaten as a food
- Avoid extracts in pregnancy

Prickly Pear – “Nopales” - *Opuntia streptacantha*

- Commonly used by Mexican people for glucose control
- Two small studies in Type II diabetes (Frati, et al. 1988; 1990)
- Single dose - 500 gm grilled or broiled stems
- Decreased fasting glucose and insulin levels
- No side effects reported
- Likely safe in pregnancy as a food



Nopales -

- May reduce blood sugar due to high fiber content
- 1 cup chopped pads = 60 calories = 5 grams fiber (20% of daily dose)
- Can be either grilled or added to vegetable dishes
- Available at many groceries

Fenugreek – *Trigonella foenum graecum*

- Legume cultivated in India, N. Africa and the Mediterranean
- Small study of Type I and II diabetics using defatted fenugreek seed powder, 100 gm/day in unleavened bread for 10 days (Sharma, et al. 1990)
- Results – decreased fasting blood sugar, postprandial glucose, and urine glucose
- Side effects - none



Fenugreek

- A study of 60 poorly controlled Type II diabetics
- Dose - 25 grams of fenugreek seed powder divided into two doses at lunch and dinner
- After 24 weeks - 47% achieved full glycemic control, 33% moderate control, 20% minimal control
- Significant improvement in fasting blood glucose and improved glucose tolerance (Sharma, et al. 1996)

Fenugreek - Safety

- Both water and alcoholic extracts exert a stimulating effect on isolated guinea pig uterus, especially during late pregnancy (Abdo MS, al Kafawi AA. 1969).
- May possess abortifacient effects, therefore, NOT recommended for use in doses higher than found in foods during pregnancy (Farnsworth et al. 1975).
- May be safe as a food
- Would NOT recommend an extract

Bottom Line for Herbs

- In pregnancy –
 - Avoid extracts
 - Food sources likely safe

Supplements in diabetes mellitus

- Chromium
- Vanadium
- Fish oil



Chromium picolinate



- Functions by regulating or potentiating insulin action
- Response time 10 days - > 3 months
- Impaired glucose tolerance appears related to chromium deficiency
- In China, 833 Type II diabetics taking 500mcg/d were followed 10 months
- Results - decreased fasting and postprandial glucose (Cheng, Zhu, Shi, 1999)
- Other studies with mixed results

Chromium picolinate

- Apparently one study in gestational diabetes
- Patients responded better to 8 mcg/kg of chromium picolinate than 4 mcg/kg (Jovanovic-Peterson, et al, 1996)
- No other studies in pregnancy
- Adequate Intake (AI) for pregnant women age 19-50 is only 30 mcg/day, above dose would be 500-600 mcg/d

Chromium picolinate - safety

- Some biochemical concern that reduction products may promote hydroxyl radicals and potential DNA damage
- However, animal studies show no carcinogenic or mutagenic effects in vivo
- Would **NOT** recommend to pregnant patients.

Vanadium



- 3 non-randomized, single-blind crossover studies (Cohen, et al.1995; Halberstam, et al. 1996; Boden, et al. 1996.)
- Total of 21 Type II diabetics
- 100 mg/d vanadyl sulfate hydrate for 3-4 weeks
- Results – decreased FBG, HgA1C, hepatic glucose production/output
- Side Effects- high rate of GI discomfort 18/21
- Insufficient evidence for safety in pregnancy for amounts > upper intake level of 1.8 mg

Fish Oil



- Cochrane Review showed no statistically significant effect on glycemic control (Farmer, et al. 2001)
- May have benefit on fetal/infant neural and retinal development when taken during last trimester of pregnancy (van Houwelingen, et al. 1995)
- May reduce risk of post partum depression
- May also be beneficial post-partum for breast-fed infants

Fish Oil

- Dose 1000 mg DHA/day
- Want a product free of methyl mercury
- Consumerlabs.com for good quality product
- Avoid fish with potentially high mercury level such as mackerel, swordfish, shark and tilefish

Yoga in Pregnancy



- Study of 166 pregnant patients - trained in yoga, relaxation, meditation and breathing exercises
- Practiced 1 hour per day
- 164 controls received routine prenatal care
- Results indicated reduced rates of prematurity, maternal HTN, and improved birth weight in intervention group (Narendran, et al. 2005)

Yoga in DM II

- Study of 24 Type II diabetics (Singh, et al. 2004)
- Sequence of 13 asanas performed 30-40 min/day for 40 days under guidance.
- Significant decrease in fasting blood glucose levels of 50 mg/dL after yoga regimen.
- The postprandial blood glucose levels decreased 75 mg/dL
- Glycosylated hemoglobin showed a decrease from 1.2 %
- Blood pressure and pulse rate improvements also noted

Mind-Body Approaches

- Relaxation techniques such as:
 - progressive muscle relaxation
 - guided imagery
 - deep breathing exercises
- May provide some benefit with regard to lowering HgA1C (Surwit, et al. 2003))
- Will benefit the fetus by reducing maternal stress hormones (Bijlani, et al. 2005)



Postpartum

- “Lifestyle behavioral changes, including weight control and exercise between pregnancies, likely would prevent recurrence of gestational diabetes as well as modify onset and severity of type 2 diabetes later in life” (Pan and associates, 1997).



Back to Mary

- Nutrition counseling and dietary changes
- May add prickly pear, beans, steel cut oats to diet for fiber, also add cinnamon
- Fish oil – DHA 1000mg/day for Mom and baby
- Exercise - Yoga
- Meditation – Relaxation techniques
- Lifestyle changes postpartum

Thank you !



Sunset in L'Anse

References

- Abdo MS, al Kafawi AA. Experimental studies on the effect of *Trigonella foenum-graecum*. *Planta Med* 1969;17(1):14-18.
- Aguwa CN, Mittal GC. Abortifacient effects of the roots of *Momordica angustisepala*. *J Ethnopharmacol* 1983;7(2):169-173.
- Baskaran K, Ahamath BK, Shanmugasundaram KR, Shanmugasundaram ERB: Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin dependent diabetes mellitus patients. *JEthnopharmacology* 30:295–305, 1990
- Baldwa VS, Bhandara CM, Pangaria A, et al. Clinical trials in patients with diabetes mellitus of an insulin-like compound obtained from plant source. *Upsala J Med Sci* 1977;82:39-41.
- Bijlani RL. Vempati RP. Yadav RK. Ray RB. Gupta V. Sharma R. Mehta N. Mahapatra SC. A brief but comprehensive lifestyle education program based on yoga reduces risk factors for cardiovascular disease and diabetes mellitus. *Journal of Alternative & Complementary Medicine*. 11(2):267-74, 2005 Apr. UI: 15865492
- Boden G, Chen X, Ruiz J, van Rossum GD, Turco S: Effects of vanadyl sulfate on carbohydrate and lipid metabolism in patients with non-insulin-dependent diabetes mellitus. *Metabolism* 45:1130–1135, 1996.

- Brankston GH, Mitchell BF, Ryan EA, et al: Resistance exercise decreases the need for insulin in overweight women with **gestational diabetes** mellitus. [Am J Obstet Gynecol 190:188, 2004](#)
- Chan WY, Tam PP, Yeung HW. The termination of early pregnancy in the mouse by beta-momorcharin. *Contraception* 1984;29(1):91-100
- Chandalia M, Garg A, Lutjohann D, et al. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. [N Engl J Med 2000;342:1392-8.](#)
- Cohen N, Halberstam M, Shlimovich P, Chang CJ, Shamoon H, Rossetti L: Oral vanadyl sulfate improves hepatic and peripheral insulin sensitivity in patients with non-insulin dependent diabetes mellitus. *J Clin Invest* 95:2501–250, 1995.
- Dempsey JC, Sorensen TK, Williams MA, et al: Prospective study of gestational diabetes mellitus in relation to maternal recreational physical activity before and during pregnancy. [Am J Epidemiol 159:663, 2004](#)
- Farmer A, Montori V, Dinneen S, Clar C. Fish oil in people with type 2 diabetes mellitus. [Review] [81 refs] Cochrane Database of Systematic Reviews. (3):CD003205, 2001. UI: 11687050

- Farnsworth NR, Bingel AS, Cordell GA, et al. Potential value of plants as sources of new antifertility agents I. J Pharm Sci 1975;64(4):535-598.
- Feig DS, Palda VA: Type 2 diabetes in pregnancy: A growing concern. [Lancet 359:1690, 2002](#)
- Frati AC, Gordillo BE, Altamirano P, Ariza CR, Cortes-Franco R, Chavez-Negrete A: Acute hypoglycemic effects of *Opuntia streptacantha* Lemiare in NIDDM (Letter). *Diabetes Care* 13:455–456, 1990
- Frati-Munari AC, Gordillo BE, Altamirano P, Ariza CR: Hypoglycemic effect of *Opuntia streptacantha* Lemaire in NIDDM. *Diabetes Care* 11:63–66, 1988
- Halberstam M, Cohen N, Shlimovich P, Rossetti L, Shamon H: Oral vanadyl sulfate improves insulin sensitivity in NIDDM but not in obese nondiabetic subjects. *Diabetes* 45:659–666, 1996.
- Helland IB, Saugstad OD, Smith L, et al. Similar effects on infants of n-3 and n-6 fatty acids supplementation to pregnant and lactating women. *Pediatrics* 2001;108(5):E82.

- Jovanovic-Peterson L, Guttierry M, Peterson CM: Chromium supplementation for gestational diabetic women (GDM) improves glucose tolerance and decreases hyperinsulinemia. *Diabetes* 43:337a, 1996.
- Johnstone FD, Nasrat AA, Prescott RJ: The effect of established and **gestational diabetes** on pregnancy outcome. [Br J Obstet Gynaecol 97:1009, 1990](#)
- Khan A, Safdar M, Ali Khan M, et al. Cinnamon improves glucose and lipids of people with type 2 diabetes. [Diabetes Care 2003;26:3215-8.](#)
- Khalsa SB. Yoga as a therapeutic intervention: a bibliometric analysis of published research studies. [Review] [181 refs] *Indian Journal of Physiology & Pharmacology*. 48(3):269-85, 2004 Jul. UI: 15648399
- Leatherdale BA, Panesar RK, Singh G, et al. Improvement in glucose tolerance due to *Momordica charantia* (karela). *Br Med J (Clin Res Ed)* 1981;282(6279):1823-1824.
- Martin JA, Hamilton BE, Sutton PD, et al: Births: Final data for 2002. *National Vital Statistics Reports*, Vol 52, No. 10. Hyattsville, MD, National Center for Health Statistics, 2003
- Narendran S. Nagarathna R. Narendran V. Gunasheela S. Nagendra HR. Efficacy of yoga on pregnancy outcome. *Journal of Alternative & Complementary Medicine*. 11(2):237-44, 2005 Apr. UI: 15865489

- Pan XR, Li GW, Hu YH, et al: Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. [Diabetes Care](#) [20:537, 1997](#)
- Pick ME, Hawrysh ZJ, Gee MI, et al. Oat bran concentrate bread products improve long-term control of diabetes: a pilot study. [J Am Diet Assoc](#) [1996;96:1254-61.](#)
- Shanmugasundaram ERB, Rajeswari G, Baskaran K, Kumar BRR, Shanmugasundaram KR, Ahmath BK: Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. *J Ethnopharmacology* 30:281–294, 1990
- Sharma RD, Raghuram TC: Hypoglycemic effect of fenugreek seeds in non-insulin dependent diabetic subjects. *Nutr Res* 10:731–739, 1990
- Sharma RD, Raghuram TC, Rao NS: Effect of fenugreek seeds on blood glucose and serum lipids in type I diabetes. *Eur J Clin Nutr* 44:301–306, 1990

- Sharma RD, Sarkar A, Hazra DK, et al. Use of fenugreek seed powder in the management of non-insulin dependent diabetes mellitus. *Nutr Res* 1996;16:1331-1339.
- Sheffield JS, Butler-Koster EL, Casey BM, et al: Maternal diabetes mellitus and infant malformations. [Obstet Gynecol 100:925, 2002](#)
- Singh S. Malhotra V. Singh KP. Madhu SV. Tandon OP. Role of yoga in modifying certain cardiovascular functions in type 2 diabetic patients. *Journal of the Association of Physicians of India*. 52:203-6, 2004 Mar.
UI: 15636309
- Surwit RS, van Tilburg MA, Zucker N, et al. Stress management improves long-term glycemic control in type 2 diabetes. *Diabetes Care* 2002;25:30-4.
- van Houwelingen AC, Sorensen JD, Hornstra G, et al. Essential fatty acid status in neonates after fish-oil supplementation during late pregnancy. *Br J Nutr* 1995;74(5):723-731.
- Yeh GY. Eisenberg DM. Kaptchuk TJ. Phillips RS. Systematic review of herbs and dietary supplements for glycemic control in diabetes. [Review] [158 refs] *Diabetes Care*. 26(4):1277-94, 2003 Apr. UI: 12663610